



# BRIGHT FUTURES HANDOUT ► PARENT

## 4 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

### ✓ HOW YOUR FAMILY IS DOING

- Learn if your home or drinking water has lead and take steps to get rid of it. Lead is toxic for everyone.
- Take time for yourself and with your partner. Spend time with family and friends.
- Choose a mature, trained, and responsible babysitter or caregiver.
- You can talk with us about your child care choices.

### ✓ FEEDING YOUR BABY

- For babies at 4 months of age, breast milk or iron-fortified formula remains the best food. Solid foods are discouraged until about 6 months of age.
- Avoid feeding your baby too much by following the baby's signs of fullness, such as
  - Leaning back
  - Turning away

#### If Breastfeeding

- Providing only breast milk for your baby for about the first 6 months after birth provides ideal nutrition. It supports the best possible growth and development.
- Be proud of yourself if you are still breastfeeding. Continue as long as you and your baby want.
- Know that babies this age go through growth spurts. They may want to breastfeed more often and that is normal.
- If you pump, be sure to store your milk properly so it stays safe for your baby. We can give you more information.
- Give your baby vitamin D drops (400 IU a day).
- Tell us if you are taking any medications, supplements, or herbal preparations.

#### If Formula Feeding

- Make sure to prepare, heat, and store the formula safely.
- Feed on demand. Expect him to eat about 30 to 32 oz daily.
- Hold your baby so you can look at each other when you feed him.
- Always hold the bottle. Never prop it.
- Don't give your baby a bottle while he is in a crib.

### ✓ YOUR CHANGING BABY

- Create routines for feeding, nap time, and bedtime.
- Calm your baby with soothing and gentle touches when she is fussy.
- Make time for quiet play.
  - Hold your baby and talk with her.
  - Read to your baby often.
- Encourage active play.
  - Offer floor gyms and colorful toys to hold.
  - Put your baby on her tummy for playtime. Don't leave her alone during tummy time or allow her to sleep on her tummy.
- Don't have a TV on in the background or use a TV or other digital media to calm your baby.

### ✓ HEALTHY TEETH

- Go to your own dentist twice yearly. It is important to keep your teeth healthy so you don't pass bacteria that cause cavities on to your baby.
- Don't share spoons with your baby or use your mouth to clean the baby's pacifier.
- Use a cold teething ring if your baby's gums are sore from teething.
- Don't put your baby in a crib with a bottle.
- Clean your baby's gums and teeth (as soon as you see the first tooth) 2 times per day with a soft cloth or soft toothbrush and a small smear of fluoride toothpaste (no more than a grain of rice).

#### Helpful Resources:

Information About Car Safety Seats: [www.safercar.gov/parents](http://www.safercar.gov/parents) | Toll-free Auto Safety Hotline: 888-327-4236

## 4 MONTH VISIT—PARENT



### SAFETY

- Use a rear-facing-only car safety seat in the back seat of all vehicles.
- Never put your baby in the front seat of a vehicle that has a passenger airbag.
- Your baby's safety depends on you. Always wear your lap and shoulder seat belt. Never drive after drinking alcohol or using drugs. Never text or use a cell phone while driving.
- Always put your baby to sleep on her back in her own crib, not in your bed.
  - Your baby should sleep in your room until she is at least 6 months of age.
  - Make sure your baby's crib or sleep surface meets the most recent safety guidelines.
  - Don't put soft objects and loose bedding such as blankets, pillows, bumper pads, and toys in the crib.
- Drop-side cribs should not be used.
- Lower the crib mattress.
- If you choose to use a mesh playpen, get one made after February 28, 2013.
- Prevent tap water burns. Set the water heater so the temperature at the faucet is at or below 120°F /49°C.
- Prevent scalds or burns. Don't drink hot drinks when holding your baby.
- Keep a hand on your baby on any surface from which she might fall and get hurt, such as a changing table, couch, or bed.
- Never leave your baby alone in bathwater, even in a bath seat or ring.
- Keep small objects, small toys, and latex balloons away from your baby.
- Don't use a baby walker.

## WHAT TO EXPECT AT YOUR BABY'S 6 MONTH VISIT

### We will talk about

- Caring for your baby, your family, and yourself
- Teaching and playing with your baby
- Brushing your baby's teeth
- Introducing solid food
- Keeping your baby safe at home, outside, and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th Edition*

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

# DTaP (Diphtheria, Tetanus, Pertussis) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

## 1. Why get vaccinated?

DTaP vaccine can prevent **diphtheria, tetanus, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **DIPHTHERIA (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

## 2. DTaP vaccine

DTaP is only for children younger than 7 years old. Different vaccines against tetanus, diphtheria, and pertussis (Tdap and Td) are available for older children, adolescents, and adults.

It is recommended that children receive 5 doses of DTaP, usually at the following ages:

- 2 months
- 4 months
- 6 months
- 15–18 months
- 4–6 years

DTaP may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

DTaP may be given at the same time as other vaccines.

## 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any **severe, life-threatening allergies**
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP or DTaP)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré Syndrome** (also called “GBS”)
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**

In some cases, your child’s health care provider may decide to postpone DTaP vaccination until a future visit.

Children with minor illnesses, such as a cold, may be vaccinated. Children who are moderately or severely ill should usually wait until they recover before getting DTaP vaccine.

Your child’s health care provider can give you more information.



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention

---

## 4. Risks of a vaccine reaction

---

- Soreness or swelling where the shot was given, fever, fussiness, feeling tired, loss of appetite, and vomiting sometimes happen after DTaP vaccination.
- More serious reactions, such as seizures, non-stop crying for 3 hours or more, or high fever (over 105°F) after DTaP vaccination happen much less often. Rarely, vaccination is followed by swelling of the entire arm or leg, especially in older children when they receive their fourth or fifth dose.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

## 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

## 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

## 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).



# Hepatitis B Vaccine:

## What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

**Hepatitis B vaccine** can prevent **hepatitis B**.

Hepatitis B is a liver disease that can cause mild illness lasting a few weeks, or it can lead to a serious, lifelong illness.

- **Acute hepatitis B infection** is a short-term illness that can lead to fever, fatigue, loss of appetite, nausea, vomiting, jaundice (yellow skin or eyes, dark urine, clay-colored bowel movements), and pain in the muscles, joints, and stomach.
- **Chronic hepatitis B infection** is a long-term illness that occurs when the hepatitis B virus remains in a person's body. Most people who go on to develop chronic hepatitis B do not have symptoms, but it is still very serious and can lead to liver damage (cirrhosis), liver cancer, and death. Chronically infected people can spread hepatitis B virus to others, even if they do not feel or look sick themselves.

Hepatitis B is spread when blood, semen, or other body fluid infected with the hepatitis B virus enters the body of a person who is not infected. People can become infected through:

- Birth (if a pregnant person has hepatitis B, their baby can become infected)
- Sharing items such as razors or toothbrushes with an infected person
- Contact with the blood or open sores of an infected person
- Sex with an infected partner
- Sharing needles, syringes, or other drug-injection equipment
- Exposure to blood from needlesticks or other sharp instruments

Most people who are vaccinated with hepatitis B vaccine are immune for life.

### 2. Hepatitis B vaccine

Hepatitis B vaccine is usually given as 2, 3, or 4 shots.

**Infants** should get their first dose of hepatitis B vaccine at birth and will usually complete the series at 6–18 months of age. **The birth dose of hepatitis B vaccine is an important part of preventing long-term illness in infants and the spread of hepatitis B in the United States.**

**Children and adolescents** younger than 19 years of age who have not yet gotten the vaccine should be vaccinated.

**Adults** who were not vaccinated previously and want to be protected against hepatitis B can also get the vaccine.

Hepatitis B vaccine is also recommended for the following people:

- People whose sex partners have hepatitis B
- Sexually active persons who are not in a long-term, monogamous relationship
- People seeking evaluation or treatment for a sexually transmitted disease
- Victims of sexual assault or abuse
- Men who have sexual contact with other men
- People who share needles, syringes, or other drug-injection equipment
- People who live with someone infected with the hepatitis B virus
- Health care and public safety workers at risk for exposure to blood or body fluids
- Residents and staff of facilities for developmentally disabled people
- People living in jail or prison
- Travelers to regions with increased rates of hepatitis B



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention



- People with chronic liver disease, kidney disease on dialysis, HIV infection, infection with hepatitis C, or diabetes

Hepatitis B vaccine may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

Hepatitis B vaccine may be given at the same time as other vaccines.

---

### 3. Talk with your health care provider

---

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of hepatitis B vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone hepatitis B vaccination until a future visit.

Pregnant or breastfeeding people should be vaccinated if they are at risk for getting hepatitis B. Pregnancy or breastfeeding are not reasons to avoid hepatitis B vaccination.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting hepatitis B vaccine.

Your health care provider can give you more information.

---

### 4. Risks of a vaccine reaction

---

- Soreness where the shot is given or fever can happen after hepatitis B vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

### 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

### 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

### 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).



# *Haemophilus influenzae* type b (Hib) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

## 1. Why get vaccinated?

Hib vaccine can prevent *Haemophilus influenzae* type b (Hib) disease.

*Haemophilus influenzae* type b can cause many different kinds of infections. These infections usually affect children under 5 years of age but can also affect adults with certain medical conditions. Hib bacteria can cause mild illness, such as ear infections or bronchitis, or they can cause severe illness, such as infections of the blood. Severe Hib infection, also called “invasive Hib disease,” requires treatment in a hospital and can sometimes result in death.

Before Hib vaccine, Hib disease was the leading cause of bacterial meningitis among children under 5 years old in the United States. Meningitis is an infection of the lining of the brain and spinal cord. It can lead to brain damage and deafness.

Hib infection can also cause:

- Pneumonia
- Severe swelling in the throat, making it hard to breathe
- Infections of the blood, joints, bones, and covering of the heart
- Death

## 2. Hib vaccine

Hib vaccine is usually given in 3 or 4 doses (depending on brand).

**Infants** will usually get their first dose of Hib vaccine at 2 months of age and will usually complete the series at 12–15 months of age.

**Children between 12 months and 5 years of age** who have not previously been completely vaccinated against Hib may need 1 or more doses of Hib vaccine.

**Children over 5 years old and adults** usually do not receive Hib vaccine, but it might be recommended for older children or adults whose spleen is damaged or has been removed, including people with sickle cell disease, before surgery to remove the spleen, or following a bone marrow transplant. Hib vaccine may also be recommended for people 5 through 18 years old with HIV.

Hib vaccine may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

Hib vaccine may be given at the same time as other vaccines.

## 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of Hib vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone Hib vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting Hib vaccine.

Your health care provider can give you more information.



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention

---

## 4. Risks of a vaccine reaction

---

- Redness, warmth, and swelling where the shot is given and fever can happen after Hib vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

## 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

## 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

## 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).





# Polio Vaccine:

## *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

**Polio vaccine** can prevent **polio**.

Polio (or poliomyelitis) is a disabling and life-threatening disease caused by poliovirus, which can infect a person's spinal cord, leading to paralysis.

Most people infected with poliovirus have no symptoms, and many recover without complications. Some people will experience sore throat, fever, tiredness, nausea, headache, or stomach pain.

A smaller group of people will develop more serious symptoms that affect the brain and spinal cord:

- Paresthesia (feeling of pins and needles in the legs),
- Meningitis (infection of the covering of the spinal cord and/or brain), or
- Paralysis (can't move parts of the body) or weakness in the arms, legs, or both.

Paralysis is the most severe symptom associated with polio because it can lead to permanent disability and death.

Improvements in limb paralysis can occur, but in some people new muscle pain and weakness may develop 15 to 40 years later. This is called "post-polio syndrome."

Polio has been eliminated from the United States, but it still occurs in other parts of the world. The best way to protect yourself and keep the United States polio-free is to maintain high immunity (protection) in the population against polio through vaccination.

### 2. Polio vaccine

**Children** should usually get 4 doses of polio vaccine at ages 2 months, 4 months, 6–18 months, and 4–6 years.

Most **adults** do not need polio vaccine because they were already vaccinated against polio as children. Some adults are at higher risk and should consider polio vaccination, including:

- People traveling to certain parts of the world
- Laboratory workers who might handle poliovirus
- Health care workers treating patients who could have polio
- Unvaccinated people whose children will be receiving oral poliovirus vaccine (for example, international adoptees or refugees)

Polio vaccine may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

Polio vaccine may be given at the same time as other vaccines.



---

### 3. Talk with your health care provider

---

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of polio vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone polio vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting polio vaccine.

Not much is known about the risks of this vaccine for pregnant or breastfeeding people. However, polio vaccine can be given if a pregnant person is at increased risk for infection and requires immediate protection.

Your health care provider can give you more information.

---

### 4. Risks of a vaccine reaction

---

- A sore spot with redness, swelling, or pain where the shot is given can happen after polio vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

### 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

### 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

### 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).



# Pneumococcal Conjugate Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

## 1. Why get vaccinated?

Pneumococcal conjugate vaccine can prevent pneumococcal disease.

**Pneumococcal disease** refers to any illness caused by pneumococcal bacteria. These bacteria can cause many types of illnesses, including pneumonia, which is an infection of the lungs. Pneumococcal bacteria are one of the most common causes of pneumonia.

Besides pneumonia, pneumococcal bacteria can also cause:

- Ear infections
- Sinus infections
- Meningitis (infection of the tissue covering the brain and spinal cord)
- Bacteremia (infection of the blood)
- Anyone can get pneumococcal disease, but children under 2 years old, people with certain medical conditions or other risk factors, and adults 65 years or older are at the highest risk.

Most pneumococcal infections are mild. However, some can result in long-term problems, such as brain damage or hearing loss. Meningitis, bacteremia, and pneumonia caused by pneumococcal disease can be fatal.

## 2. Pneumococcal conjugate vaccine

Pneumococcal conjugate vaccine helps protect against bacteria that cause pneumococcal disease. There are three pneumococcal conjugate vaccines (PCV13, PCV15, and PCV20). The different vaccines are recommended for different people based on their age and medical status.

### PCV13

- **Infants and young children** usually need 4 doses of PCV13, at ages 2, 4, 6, and 12–15 months.
- **Older children (through age 59 months)** may be vaccinated with PCV13 if they did not receive the recommended doses.
- **Children and adolescents 6–18 years of age** with certain medical conditions should receive a single dose of PCV13 if they did not already receive PCV13.

### PCV15 or PCV20

- **Adults 19 through 64 years old** with certain medical conditions or other risk factors who have not already received a pneumococcal conjugate vaccine should receive either:
  - a single dose of PCV15 followed by a dose of pneumococcal polysaccharide vaccine (PPSV23), or
  - a single dose of PCV20.
- **Adults 65 years or older** who have not already received a pneumococcal conjugate vaccine should receive either:
  - a single dose of PCV15 followed by a dose of PPSV23, or
  - a single dose of PCV20.

Your health care provider can give you more information.



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention

---

### 3. Talk with your health care provider

---

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any type of pneumococcal conjugate vaccine (PCV13, PCV15, PCV20, or an earlier pneumococcal conjugate vaccine known as PCV7), or to any vaccine containing diphtheria toxoid (for example, DTaP), or has any severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone pneumococcal conjugate vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover.

Your health care provider can give you more information.

---

### 4. Risks of a vaccine reaction

---

- Redness, swelling, pain, or tenderness where the shot is given, and fever, loss of appetite, fussiness (irritability), feeling tired, headache, muscle aches, joint pain, and chills can happen after pneumococcal conjugate vaccination.

Young children may be at increased risk for seizures caused by fever after PCV13 if it is administered at the same time as inactivated influenza vaccine. Ask your health care provider for more information.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

### 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**.

*VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

### 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

### 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).



# Rotavirus Vaccine:

## *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

Rotavirus vaccine can prevent rotavirus disease.

Rotavirus commonly causes severe, watery diarrhea, mostly in babies and young children. Vomiting and fever are also common in babies with rotavirus. Children may become dehydrated and need to be hospitalized and can even die.

### 2. Rotavirus vaccine

Rotavirus vaccine is administered by putting drops in the child's mouth. Babies should get 2 or 3 doses of rotavirus vaccine, depending on the brand of vaccine used.

- The first dose must be administered before 15 weeks of age.
- The last dose must be administered by 8 months of age.

Almost all babies who get rotavirus vaccine will be protected from severe rotavirus diarrhea.

Another virus called "porcine circovirus" can be found in one brand of rotavirus vaccine (Rotarix). This virus does not infect people, and there is no known safety risk.

Rotavirus vaccine may be given at the same time as other vaccines.

### 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of rotavirus vaccine**, or has any **severe, life-threatening allergies**
- Has a **weakened immune system**
- Has **severe combined immunodeficiency (SCID)**
- Has had a type of bowel blockage called "**intussusception**"

In some cases, your child's health care provider may decide to postpone rotavirus vaccination until a future visit.

Infants with minor illnesses, such as a cold, may be vaccinated. Infants who are moderately or severely ill should usually wait until they recover before getting rotavirus vaccine.

Your child's health care provider can give you more information.

### 4. Risks of a vaccine reaction

- Irritability or mild, temporary diarrhea or vomiting can happen after rotavirus vaccine.

Intussusception is a type of bowel blockage that is treated in a hospital and could require surgery. It happens naturally in some infants every year in the United States, and usually there is no known reason for it. There is also a small risk of intussusception from rotavirus vaccination, usually within a week after the first or second vaccine dose. This additional risk is estimated to range from about 1 in 20,000 U.S. infants to 1 in 100,000 U.S. infants who get rotavirus vaccine. Your health care provider can give you more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention



---

## 5. What if there is a serious problem?

---

For intussusception, look for signs of stomach pain along with severe crying. Early on, these episodes could last just a few minutes and come and go several times in an hour. Babies might pull their legs up to their chest. Your baby might also vomit several times or have blood in the stool, or could appear weak or very irritable. These signs would usually happen during the first week after the first or second dose of rotavirus vaccine, but look for them any time after vaccination. If you think your baby has intussusception, contact a health care provider right away. If you can't reach your health care provider, take your baby to a hospital. Tell them when your baby got rotavirus vaccine.

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

## 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

## 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).

